

Introduction to Group Theory

Note 4 Group Algebra

August 10, 2009

Contents

1	SU(n) Algebra	1
2	SU(2) Algebra	2
3	Representation of SU(2) algebra	2
4	Product Representation	4
5	SU(3) Algebra	6
6	Representation of SU(3)	7

Algebraic Approach

For groups larger than $SU(2)$, it is more efficient to take the algebraic approach. This is the study of the structure of group in terms of commutation relation of the generators, similar to the study of rotation group in terms of the angular momentum algebra in quantum mechanics. In fact, the angular momentum algebra forms the basis for the irreducible representations of more general groups.

1 SU(n) Algebra

We will now study the $SU(2)$ in terms of their generators. More generally $SU(n)$ is the group formed by $n \times n$ unitary matrices with unit determinant. We now compute the number of independent real group parameters needed to represent the group elements. The unitary condition $UU^\dagger = 1$, implies

$$\sum_k U_{ik}U_{jk}^* = \delta_{ij}$$

Define n -dimensional vectors by

$$\vec{U}_i = (U_{i1}, U_{i2}, \dots, U_{in}), \quad i = 1, 2, 3, \dots, n$$

Then unitary conditions can be written as

$$\left(\vec{U}_i, \vec{U}_j\right) = \sum_k U_{ik}U_{jk}^* = \delta_{ij}$$

This means \vec{U}_i 's form an orthonormal set and

$$\left(\vec{U}_i, \vec{U}_i\right) = 1$$

gives n conditions on the real components of matrix elements and

$$\left(\vec{U}_i, \vec{U}_j\right) = 0 \quad \text{for } i \neq j \text{ give } \frac{n(n-1)}{2} \times 2 = n^2 - n \quad \text{conditions}$$

$$\det U = 1 \quad \text{gives one condition}$$

Hence, the number of independent real parameters in $SU(n)$ is

$$2n^2 - (n^2 - n) - n - 1 = n^2 - 1$$

Another way to get the same result is use the fact that any unitary matrix U can be written in terms of hermitian matrix H as

$$U = e^{iH}$$

Using the identity

$$\det(e^A) = e^{\text{Tr}A}$$

we can see that

$$\det U = 1, \quad \text{Tr}H = 0$$

It is easy to see that in terms of $n \times n$ traceless hermitian matrices, $SU(n)$ has $n^2 - 1$ parameters.

2 $SU(2)$ Algebra

We now study the case of $SU(2)$. Physical examples of $SU(2)$ group are the rotation group in 3-dimension and isospin symmetry in the strong interaction. As we have discussed before, there are 3 parameters for this group. Parametrize 2×2 unitary matrices by

$$U(\varepsilon_1, \varepsilon_2, \varepsilon_3) = \exp(i\varepsilon_i \sigma_i)$$

where $\varepsilon_1, \varepsilon_2, \varepsilon_3$ are real group parameters, and $\sigma_1, \sigma_2, \sigma_3$ are the usual Pauli matrices. The commutation relations for Pauli matrices are of the form,

$$\left[\frac{\sigma_i}{2}, \frac{\sigma_j}{2} \right] = i\varepsilon_{ijk} \frac{\sigma_k}{2}, \quad i, j, k = 1, 2, 3$$

The operators J_1, J_2, J_3 which satisfy the same commutation relation

$$[J_i, J_j] = i\varepsilon_{ijk} J_k \tag{1}$$

are called the generators of $SU(2)$ and the commutation relations are called the **Lie algebra of $SU(2)$** . Clearly these commutation relations are the same as those of the angular momentum operators $\vec{L} = \vec{r} \times \vec{p}$,

$$[L_i, L_j] = i\varepsilon_{ijk} L_k \tag{2}$$

Thus these two algebras (1)(2) are in one to one correspondance(isomorphism). Here we have set $\hbar = 1$ for convenience.

Recall that the angular momentum operators are given by

$$L_1 = -i \left(x_2 \frac{\partial}{\partial x_3} - x_3 \frac{\partial}{\partial x_2} \right), \quad L_2 = -i \left(x_3 \frac{\partial}{\partial x_1} - x_1 \frac{\partial}{\partial x_3} \right), \quad L_3 = -i \left(x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1} \right)$$

It is clear that these operators will leave the quadratic form,

$$x_1^2 + x_2^2 + x_3^2$$

invariant.

3 Representation of $SU(2)$ algebra

Any set of matrices D_1, D_2, D_3 which satisfy the same algebra,

$$[D_i, D_j] = i\varepsilon_{ijk} D_k$$

are called the **representation** of generators J_1, J_2, J_3 . For example $D_i = \frac{\sigma_i}{2}$ is a representation (defining representation or fundamental representation)

To find all other representations, we define

$$J_{\pm} = J_1 \pm iJ_2$$

Then we get

$$[J_{\pm}, J_3] = \mp J_{\pm}, \quad [J_+, J_-] = 2J_3$$

Suppose $|m\rangle$ is an eigenstate of J_3 with eigenvalue m ,

$$J_3 |m\rangle = m |m\rangle$$

then we get from these commutation relations,

$$J_3 (J_+ |m\rangle) = (m+1) (J_+ |m\rangle), \quad J_3 (J_- |m\rangle) = (m-1) (J_- |m\rangle)$$

Thus J_+ (J_-) is the raising(lowering) operator which increases(decrease) the eigenvalues by one unit. Define the total angular momentum operator by,

$$J^2 \equiv J_1^2 + J_2^2 + J_3^2 = \frac{1}{2} (J_+ J_- + J_- J_+ + J_3^2) \geq 0 \quad (3)$$

Then we get

$$[J^2, J_i] = 0, \quad \text{for } i = 1, 2, 3$$

and J^2 is called **Casmir operator**, the operator which commutes with all the generators in the group. From Schur's lemma we know that the Casmir operators will be represented by a multiple of identity matrix in any irrep. Choose the states to be eigenstates of J^2, J_3 , with eigenvalues, λ, m

$$J^2 |\lambda, m\rangle = \lambda |\lambda, m\rangle, \quad J_3 |\lambda, m\rangle = m |\lambda, m\rangle$$

with normalization

$$\langle \lambda', m' | \lambda, m \rangle = \delta_{\lambda\lambda'} \delta_{mm'} \quad (4)$$

Note that from Eq(3) we see that m , the eigenvalue of J_3 is bounded by

$$m^2 \leq \lambda$$

This will make the representation matrices finite dimensional. Since $J_{\pm} |\lambda, m\rangle$ are eigenstates of J_3 with eigenvalues $m \pm 1$, we can write

$$J_{\pm} |\lambda, m\rangle = C_{\pm}(\lambda, m) |\lambda, m \pm 1\rangle,$$

Here $C_{\pm}(\lambda, m)$ are constants to be determined by the normalization conditions in Eq(4). Since $\lambda - m^2 \geq 0$, eigenvalue m^2 is bounded. This means that for largest value of m , say $m = j$, we have

$$J_+ |\lambda, j\rangle = 0,$$

We can write J^2 operator as

$$J^2 = \frac{1}{2} (J_+ J_- + J_- J_+ + J_3^2) = J_- J_+ + J_3^2 + J_3$$

Applying this on the state $|\lambda, j\rangle$, we get

$$(\lambda - j^2 - j) |\lambda, j\rangle = 0, \quad \Rightarrow \quad \lambda = j(j+1)$$

Similarly, for smallest value of m , say $m = j'$, we have

$$J_- |\lambda, j'\rangle = 0, \quad \text{and} \quad \lambda = j'(j'-1)$$

Combining these two cases, we get

$$j(j+1) = j'(j'-1), \quad \Rightarrow \quad j = -j', \quad \text{or} \quad j' = j+1$$

The solution $j' = j+1$ violates the assumption that j is the largest value of m . Thus we will take $j = -j'$. Since J_- decreases value of m by 1 each time it acts on the eigenstate of J_3 , this implies that the maximum and minimum values of m should differ by an integer,

$$j - j' = 2j = \text{integer}, \quad \Rightarrow \quad j \text{ integer or half integer}$$

We will now use the parameter j to label the state instead of λ , with

$$J^2 |j, m\rangle = j(j+1) |j, m\rangle$$

The coefficients $C_{\pm}(\lambda, m)$ can be calculated as follows,

$$J_+ |j, m\rangle = C_+(j, m) |j, m+1\rangle, \quad \langle j, m | J_- = \langle j, m+1 | C_+^*(j, m)$$

$$\langle j, m | J_- J_+ | j, m \rangle = |C_+(j, m)|^2$$

On the other hand,

$$\langle j, m | J_- J_+ | j, m \rangle = \langle j, m | (J^2 - J_3^2 - J_3) | j, m \rangle = [j(j+1) - m^2 - m] = (j-m)(j+m+1)$$

We can then take

$$C_+(j, m) = \sqrt{(j-m)(j+m+1)}$$

Similarly,

$$C_-(j, m) = \sqrt{(j+m)(j-m+1)}$$

To summarize, the states $|j, m\rangle$, $m = -j, -j+1, \dots, j-1, j$ form the basis of the irreducible representation characterized by j . These states have the following properties,

$$J^2 |j, m\rangle = j(j+1) |j, m\rangle, \quad J_3 |j, m\rangle = m |j, m\rangle \quad (5)$$

$$J_+ |j, m\rangle = \sqrt{(j-m)(j+m+1)} |j, m+1\rangle, \quad J_- |j, m\rangle = \sqrt{(j+m)(j-m+1)} |j, m-1\rangle \quad (6)$$

Note that we can get the matrix elements of J_1, J_2 by using the relations,

$$J_1 = \frac{1}{2}(J_+ + J_-), \quad J_2 = \frac{1}{2i}(J_+ - J_-)$$

Thus for a given value of j , the matrices constructed for J_1, J_2 and J_3 will satisfy the angular momentum algebra given in Eq(1) and they are the irreps of $SU(2)$ group.

Remark:

For the case of Lorentz group in 2 dimensions, the generators are,

$$L_3 = -i \left(x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1} \right), \quad K_1 = -i \left(x_0 \frac{\partial}{\partial x_1} + x_1 \frac{\partial}{\partial x_0} \right), \quad K_2 = -i \left(x_0 \frac{\partial}{\partial x_2} + x_2 \frac{\partial}{\partial x_0} \right)$$

These are the operators which leave the quadratic form,

$$x_1^2 + x_2^2 - x_0^2$$

invariant. From these generators we get the commutation relations,

$$[L_3, K_1] = iK_2, \quad [L_3, K_2] = -iK_1, \quad [K_1, K_2] = -iL_3$$

We can also define the raising and lowering operators by

$$K_+ = K_1 + iK_2, \quad K_- = K_1 - iK_2$$

to get the commutation relations,

$$[L_3, K_\pm] = \mp K_\pm, \quad [K_+, K_-] = 2L_3$$

We also get that

$$[K_1^2 + K_2^2 - L_3^2, L_3] = 0, \quad [K_1^2 + K_2^2 - L_3^2, K_1] = 0, \quad [K_1^2 + K_2^2 - L_3^2, K_2] = 0$$

The Casimir operator is now of the form $B = K_1^2 + K_2^2 - L_3^2$. In this case the eigenvalue of L_3 is no longer bounded by that of B . The main difference between Lorentz group in 2 dimension and the 3-dimensional rotation group $O(3)$ is that $O(3)$ is a compact group while the Lorentz group is non-compact.

4 Product Representation

In the physical application of $SU(2)$ group, we need to deal with product representations. For example, if we have 2 spin 1/2 particles, we want to know the total spin J of the product of the two wavefunctions. In this simple case, the answer is $J = 0$, or 1. We now want to study this problem in terms of group theory. Denote the spin-up and spin-down states of the first particle by r_1 and r_2 , and for the second particles, s_1 and s_2 . Under $SU(2)$ matrices, they transform according to

$$r'_i = U \left(\vec{\epsilon} \right)_{ij} r_j, \quad s'_k = U \left(\vec{\epsilon} \right)_{kl} s_l$$

where

$$U \left(\vec{\epsilon} \right) = \exp \left(i \vec{\epsilon} \cdot \vec{J} \right), \quad \text{and} \quad \vec{J} = \frac{\vec{\sigma}}{2}$$

Then the product will transform as

$$(r'_i s'_k) = U\left(\vec{\epsilon}\right)_{ij} U\left(\vec{\epsilon}\right)_{kl} (r_j s_l) = D\left(\vec{\epsilon}\right)_{ik,jl} (r_j s_l)$$

where

$$D\left(\vec{\epsilon}\right)_{ik,jl} = U\left(\vec{\epsilon}\right)_{ij} U\left(\vec{\epsilon}\right)_{kl}$$

Generally, $D\left(\vec{\epsilon}\right)$ is reducible. To see what irreducible representation it decomposes into, it is easier to work with the generators directly by taking $\varepsilon_i \ll 1$,

$$r'_i \simeq \left(1 + i\vec{\epsilon} \cdot \vec{J}\right)_{ij} r_j = \left(1 + i\vec{\epsilon} \cdot \vec{J}^{(1)}\right)_{ij} r_j$$

$$s'_k \simeq \left(1 + i\vec{\epsilon} \cdot \vec{J}\right)_{kl} s_l = \left(1 + i\vec{\epsilon} \cdot \vec{J}^{(2)}\right)_{kl} s_l$$

where $\vec{J}^{(1)}$ is defined to operate on r_i and does not affect s_k ; while $\vec{J}^{(2)}$ operates on s_i and does not affect r_k . Define the total angular momentum operator as

$$\vec{J} = \vec{J}^{(1)} + \vec{J}^{(2)}$$

We now change to more familiar notation. Let α_i denote the spin-up state of the i th particle and β_i the spin-down state of the i th particle. There are four combinations of two-particle states : $\alpha_1\alpha_2, \alpha_1\beta_2, \beta_1\alpha_2, \beta_1\beta_2$. We start with state with largest value of J_3 ,

$$J_3 |\alpha_1\alpha_2\rangle = J_3^{(1)} |\alpha_1\alpha_2\rangle + J_3^{(2)} |\alpha_1\alpha_2\rangle = |\alpha_1\alpha_2\rangle$$

To find the total angular momentum, we write

$$\left(\vec{J}\right)^2 = \left(\vec{J}^{(1)}\right)^2 + \left(\vec{J}^{(2)}\right)^2 + \left[\left(J_+^{(1)} J_-^{(2)} + J_-^{(1)} J_+^{(2)}\right) + 2J_3^{(1)} J_3^{(2)}\right]$$

and find that

$$\left(\vec{J}\right)^2 |\alpha_1\alpha_2\rangle = 2 |\alpha_1\alpha_2\rangle$$

This means the state $|\alpha_1\alpha_2\rangle$ has $J = 1$ and $J_3 = 1$,

$$|1, 1\rangle = |\alpha_1\alpha_2\rangle \tag{7}$$

We can use the lowering operator $J_- = J_-^{(1)} + J_-^{(2)}$, to reach all other states in the $J = 1$ irrep,

$$J_- |1, 1\rangle = J_- |\alpha_1\alpha_2\rangle = \left(J_-^{(1)} + J_-^{(2)}\right) |\alpha_1\alpha_2\rangle = |\alpha_1\beta_2\rangle + |\beta_1\alpha_2\rangle$$

On the other hand, from Eq(6) we get,

$$J_- |1, 1\rangle = \sqrt{2} |1, 0\rangle$$

Then

$$|1, 0\rangle = \frac{1}{\sqrt{2}} (|\alpha_1\beta_2\rangle + |\beta_1\alpha_2\rangle) \tag{8}$$

Obviously,

$$|1, -1\rangle = |\beta_1\beta_2\rangle$$

The remaining independent state with $J = 0$ can be obtained by orthogonality with respect to the state $|1, 0\rangle$ given in Eq(8),

$$|0, 0\rangle = \frac{1}{\sqrt{2}} (|\alpha_1\beta_2\rangle - |\beta_1\alpha_2\rangle)$$

More generally, the product representations, $|j_1, m_1\rangle \times |j_2, m_2\rangle$ can be combined into eigenstates $|J, M\rangle$ of total angular momentum, $\vec{J} = \vec{J}^{(1)} + \vec{J}^{(2)}$,

$$|J, M\rangle = \sum_{m_1, m_2} |j_1, m_1\rangle |j_2, m_2\rangle \langle j_1, m_1, j_2, m_2 | JM \rangle$$

The coefficients $\langle j_1, m_1, j_2, m_2 | J, M \rangle$ are called **Clebsch-Gordon coefficients**. Thus for the above case (Eqs(7,8))

$$\left\langle \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2} | 1, 1 \right\rangle = 1, \quad \left\langle \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2} | 1, 0 \right\rangle = \frac{1}{\sqrt{2}}, \quad \text{etc}$$

Note that the J_3 quantum number is additive,

$$M = m_1 + m_2$$

The procedure of working out the irrep of the product representations can be summarized as follows.

1. Start with the combination of states with largest J_3 . Clearly this is also an eigenstate with largest total J .
2. Use the lowering operator $J_- = J_-^{(1)} + J_-^{(2)}$ to get all the other states in the same irrep.
3. Find the orthogonal combination to $|J_m, J_m - 1\rangle$ where J_m is the maximum value of J in the product. This orthogonal state should be $|J_m - 1, J_m - 1\rangle$. Then use the lowering operator to reach other $J = J_m - 1$ states.
4. Repeat these steps until $J = |j_1 - j_2|$.

5 $SU(3)$ Algebra

Physical example of this group is the eightfold way of mesons and baryons. Moreover this forms the basis for the quark model in particle physics. Denote the group parameters by α_i , $i = 1, 2, \dots, 8$ and write the group elements as

$$U(\alpha_i) = \exp(i\alpha_i \lambda_i), \quad \lambda_i : \text{hermitian traceless } 3 \times 3 \text{ matrices}$$

The standard form of λ_i are those first given by Gell-Mann,

$$\begin{aligned} \lambda_1 &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \lambda_4 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \quad \lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \\ \lambda_8 &= \sqrt{\frac{1}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \end{aligned}$$

Note that these matrices are normalized as

$$\text{Tr}(\lambda_i \lambda_j) = 2\delta_{ij}$$

The commutators for these matrices, the Lie algebras, are of the form,

$$\left[\frac{\lambda_i}{2}, \frac{\lambda_j}{2} \right] = if_{ijk} \left(\frac{\lambda_k}{2} \right),$$

where f_{ijk} are the totally anti-symmetric structure constants and those which are non-zero are given by,

$$f_{123} = 1, \quad f_{147} = -f_{156} = f_{246} = f_{257} = f_{345} = -f_{367} = \frac{1}{2}, \quad f_{458} = f_{678} = \frac{\sqrt{3}}{2}$$

Thus the generators F_i of $SU(3)$ satisfy the same commutators,

$$[F_i, F_j] = if_{ijk} F_k$$

From the fact that λ_3 and λ_8 are diagonal, we have

$$[\lambda_3, \lambda_8] = 0$$

which implies that

$$[F_3, F_8] = 0$$

Hence, F_3 and F_8 can be diagonalized simultaneously and their eigenvalues, up to some scaling factor, will be used to label states in the representation. The largest number of mutually commuting generators in the algebra is called the **rank of the algebra**. Thus $SU(3)$ is a rank 2 group while $SU(2)$ is a rank 1 group. This is also the number of eigenvalues needed to label the states in a given irreps.

6 Representation of $SU(3)$

Define the raising and lowering operators by

$$T_{\pm} = F_1 \pm iF_2, \quad U_{\pm} = F_6 \pm iF_7, \quad V_{\pm} = F_4 \pm iF_5$$


If we write $T_3 = F_3$, $Y = \frac{2}{\sqrt{3}}F_8$, where T_3 is the usual isospin and Y is the hypercharge, the commutation relations become

$$\begin{aligned} [T_3, T_{\pm}] &= \pm T_{\pm} & [Y, T_{\pm}] &= 0 & [T_+, T_-] &= 2T_3 \\ [T_3, U_{\pm}] &= \mp \frac{1}{2}U_{\pm} & [Y, U_{\pm}] &= \pm U_{\pm} & [U_+, U_-] &= \frac{3}{2}Y - T_3 \equiv 2U_3 \\ [T_3, V_{\pm}] &= \pm \frac{1}{2}V_{\pm} & [Y, V_{\pm}] &= \pm V_{\pm} & [V_+, V_-] &= \frac{3}{2}Y + T_3 \equiv 2V_3 \\ [T_+, V_+] &= 0 & [T_+, U_-] &= 0 & [U_+, V_+] &= 0 \\ [T_+, V_-] &= -U_- & [T_+, U_+] &= V_+ & [U_+, V_-] &= T_- \end{aligned}$$


Clearly, these raising and lowering operators move the states on the plane labeled by (T_3, Y) ,

$$\begin{aligned} T_+ & \text{ raises } T_3 \text{ by 1 unit and leaves } y \text{ unchanged;} \\ U_+ & \text{ raises } T_3 \text{ by } \frac{1}{2} \text{ unit and raises } y \text{ by 1 unit;} \\ V_+ & \text{ raises } T_3 \text{ by } \frac{1}{2} \text{ unit and raises } y \text{ by 1 unit;} \text{ etc } \dots \end{aligned}$$

If the units of T_3 and Y are appropriately scaled in the graph, these raising and lowering operators connect points along lines that are multiple of 60° with respect to each other.



Each irrep of $SU(3)$ is characterized by a set of two integers (p, q) . Graphically it shows up as a figure with a hexagonal boundary on the $T_3 - Y$ plane: three sides having p units of length and other sides having q units: the hexagon collapses into an equilateral triangle when either p or q vanishes. The boundary is symmetric under reflection about Y -axis. We recall that an $SU(2)$ irreps is characterized by one integer j ; graphically it is a straight line of $2j$ length. There are $2j + 1$ sites, each of them singly occupied by one state. For the $SU(3)$ representation (p, q) the multiplicity of states on each site in the $T_3 - Y$ plane form the following pattern: the sites in the boundary are singly occupied, on the next layer they are doubly occupied, on the third layer they are triply occupied, etc., until a triangular layer is reached beyond which the multiplicity ceases to increase and remains to be $q + 1$ for $p > q$ ($p + 1$ for $q > p$).



To get all the states within a given irrep, start with state with largest value of T_3 , ψ_{\max} , which can be characterized as

$$T_+ \psi_{\max} = V_+ \psi_{\max} = U_- \psi_{\max} = 0$$

Repeatedly apply V_- on ψ_{\max} , it has to vanish at some point, say after $p + 1$ steps,

$$(V_-)^{p+1} \psi_{\max} = 0$$

At the step $(V_-)^p \psi_{\max}$, we can apply T_- until it reach another corner,

$$(T_-)^{q+1} (V_-)^p \psi_{\max} = 0$$

Then we use U_+, V_+, T_+, U_- until it comes back ψ_{\max} . This completes the outer layer of the representation. For the next layer we can start with state ψ'_{\max} which has same properties as ψ_{\max} , *i.e.*

$$T_+ \psi'_{\max} = V_+ \psi'_{\max} = U_- \psi'_{\max} = 0$$

but with T_3 quantum number one unit less than ψ_{\max} . Repeat the procedures as described above.

Remarks :

1. The boundary is always convex.

This can be seen as follows. If there is a state at site C as shown in the figure, we can write


$$U_+ \psi_C = \lambda \psi_A, \quad \lambda \text{ some constant and } \psi_A = V_- \psi_B$$

Then

$$\lambda (\psi_A, \psi_A) = (V_- \psi_B, U_+ \psi_C) = (\psi_B, V_+ U_+ \psi_C) = (\psi_B, U_+ V_+ \psi_C) = 0$$

This implies $\lambda = 0$ and there is no state at site C .

2. The boundary will in general be a six-sided figure and is symmetric under rotation of 120° . The boundary is also symmetric with respect to reflection in $Y - axis$.
3. There is one state at each site on the boundary, 2 states at each site on the next layer, 3 states at each site on the next layer, etc, until a triangular layer is reached, beyond which the multiplicity ceases to increase. Denote the representation by the integers (p, q)



Then the number of states in the inner most triangle is

$$\sum_{l=1}^{p-q+1} l = \frac{1}{2} (p - q + 1) (p - q + 2)$$

and the multiplicity for each state in the triangle is $(q + 1)$. The number of states in the next layer is $3(p - q + 2)$ with multiplicity q . Continuing the counting of states in this fashion, we can get the dimensionality of the irrep as

$$\begin{aligned} d(p, q) &= \frac{1}{2} (q + 1) (p - q + 1) (p + q + 2) + 3 \sum_{\nu=0}^q (q - \nu) (p - q + 2\nu + 2) \\ &= \frac{1}{2} (p + 1) (q + 1) (p + q + 2) \end{aligned}$$

4. In general, there will be several states for a given value of (T_3, Y) . The states on the $Y = \text{constant}$ line form isospin multiplet. For example, there are $p + 1$ states on the top line and the isospin of these states is $T = \frac{p}{2}$. Similarly the next line will have isospin $T = \frac{1}{2} (p - 1)$, etc. It is not hard to see that $(T_3)_{\max} = \frac{1}{2} (p + q)$

