

Dark photon jets

Presented at Fo-Guang-Shan

Hsiang-nan Li

Dec. 29th, 2018

Collaborated with J.M. Chen, P. Ko, J.M. Li, H. Yokoya

Motivation

- BSM physics should exist
- Dark matter expected, despite of null results from direct and indirect detections so far
- Given rich dynamics in SM, natural to think about interactive structure of dark sector
- Dark sector can be explored in QCD aspect through its very weak coupling to SM

Dark photon jets

- Possible to have hidden $U(1)'$ gauge group---dark photons
- $U(1)'$ may kinetically mix with SM $U(1)$
- Light (sub-GeV) DM charged under $U(1)'$, if produced energetically at collider, radiates collimated dark photons, decaying back to SM particles (leptons, hadrons), and forming jets
- Jet substructures (intensively studied in QCD) of dark photons can reveal DM property

Chirality and mass generation

- Here determine DM fermion is chiral- or vector-like by dark photon jet substructure
- Chirality of DM fermion might be related to mass generation mechanism
- Particle mass usually generated by Higgs mechanism
- For $U(1)$ gauge group, Stueckelberg mechanism is also possible


Higgs mechanism 1964

- Before symmetry breaking

$$\mathcal{L} = -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + \frac{\epsilon}{2}F'_{\mu\nu}F^{\mu\nu} + |D_\mu\Phi'|^2 - \frac{\lambda_{\Phi'}}{4}\left(|\Phi'|^2 - \frac{v_{\Phi'}^2}{2}\right)^2$$

$$+ \sum_{s=L/R} i\bar{\chi}_s \not{D} \chi_s - (y_\chi \bar{\chi}_L \Phi' \chi_R + h.c.),$$

$D_\mu = \partial_\mu - ig'Q_s A'_\mu \quad s = L/R \quad Q_{\Phi'} = Q_L - Q_R$



- After symmetry breaking

$$\Phi' = \frac{1}{\sqrt{2}}(h' + i\phi') \quad h' \rightarrow h' + \overset{\text{vev}}{v_{\Phi'}}$$

$$m_{A'} = g'Q_{\Phi'}v_{\Phi'} \quad m_\chi = \frac{y_\chi v_{\Phi'}}{\sqrt{2}} \quad m_{h'}^2 = \frac{\lambda_{\Phi'} v_{\Phi'}^2}{2}$$

Stueckelberg mechanism 1938

- Limit of Higgs model with vev going to infinity and Higgs charge, Yukawa coupling to zero in a way that gauge boson and fermion masses stay fixed
- Higgs with infinite mass decouples
- Theory remains renormalizable though not manifestly gauge invariant

$$\mathcal{L} = -\frac{1}{4}F'_{\mu\nu}F'^{\mu\nu} + \frac{\epsilon}{2}F'_{\mu\nu}F^{\mu\nu} + \frac{1}{2}m_{A'}^2 A'_\mu A'^\mu + \sum_s \bar{\chi}_s (i\not{D} - m_\chi) \chi_s$$

$Q_L = Q_R$ Higgs charge goes to zero

Higgs vs Stueckelberg

- If DM fermion chiral-like, left- and right-handed fermions can have different $U(1)'$ charges. Bare fermion mass term is forbidden
- Dark Higgs exists to give DM fermion and dark photon masses
- If DM fermion vector-like, left- and right-handed fermions have same charge
- **Naturally** assume dark photon mass comes from Stueckelberg mechanism. No Higgs

Models and Parameters

- Chiral (Higgs) vs vector model (Stueckelberg)
- Difference characterized by charge ratio Q_L/Q_R

$$(Q_L, Q_R) = (2, 0) \quad \text{for the Chiral Model}$$

$$(Q_L, Q_R) = (1, 1) \quad \text{for the Vector Model}$$

- Benchmark points


point A: $\alpha' = 0.3$ $m_\chi = 0.7$ GeV $m_{A'} = 0.4$ GeV $m_{h'} = 1.0$ GeV,

point B: $\alpha' = 0.15$ $m_\chi = 1.0$ GeV $m_{A'} = 0.4$ GeV $m_{h'} = 1.0$ GeV,

point C: $\alpha' = 0.075$ $m_\chi = 1.4$ GeV $m_{A'} = 0.4$ GeV $m_{h'} = 1.4$ GeV

Dark shower

dark photon with
transverse polarization



dark Higgs decays into
dark photon pair

Splitting functions

Chen, Han, Tweedie 2016

$$\frac{d\mathcal{P}_{A \rightarrow B+C}}{dz dk_T^2} \quad z = \frac{E_C}{E_A}$$

soft singularity

$$\frac{d\mathcal{P}}{dz dk_T^2}(\chi_s \rightarrow \chi_s + A'_T) = \frac{\alpha'}{2\pi} Q_s^2 \frac{1 + \bar{z}^2}{z} \frac{k_T^2}{\tilde{k}_T^4},$$

$\leftarrow k_T^2 + \bar{z}^2 m_\chi^2 + z m_{A'}^2$

$$\frac{d\mathcal{P}}{dz dk_T^2}(\chi_s \rightarrow \chi_{-s} + A'_L) = \frac{\alpha'}{2\pi} \frac{m_\chi^2}{m_{A'}^2} Q_{\Phi'}^2 \frac{z}{2} \frac{k_T^2}{\tilde{k}_T^4}$$

vanish for
vector model

$$\frac{d\mathcal{P}}{dz dk_T^2}(\chi_s \rightarrow \chi_{-s} + h')$$

helicity flip, proportional to fermion mass

Setting

- DM fermion pair production at LHC with c.o.m $E=14$ TeV through effective operator $(\bar{q}\gamma^\mu q)(\bar{\chi}\gamma_\mu\chi)$
- Plus associated jet with $p_T > 200$ GeV to have missing energy
- Total width $\Gamma_{A'} \sim \alpha_{\text{em}}\epsilon^2 M_{A'}$ corresponding to A' decay length $\mathcal{O}(1)$ mm demands large enough kinetic mixing $\epsilon \gtrsim 8.2 \times 10^{-6}$, so that dark photons mostly decay into SM particles inside collider

Observables


- Implement splitting functions into dark shower
- Final-state dark radiation only, since initial state SM radiation mainly soft (of order GeV) with jet p_T cut (of order 100 GeV), and negligible
- Consider IR safe observables, like scalar sum:

$$H_T = \sum_{i=A'} |p_{T_i}|$$


- Number of dark photons $n_{A'}$ is not IR safe at high energy Zhang, Kim, Lee, Park 2016
- Dark photon Jet substructures

Clustering dark photons


- Anti kT algorithm for radius $R=2$ to determine jet axis
- Average energy deposit over 10^4 DM jet events
- Find jet profile $f_E(r)$, defined as energy fraction outside cone of radius $r < R$





Chiral model gives wider jets




Ideal observable




Results for HT





difference between chiral
and vector enhanced by
Yukawa coupling $\frac{m_\chi}{m_{A'}}$





confirm that almost all dark photons decay into SM particles

confirm that HT is IR safe, but dark photon number is not



Summary

- Jet substructures useful for revealing properties of parent particles
- Dark sector may have interactive behavior, and $U(1)'$ interaction is a simple scenario
- Dark photon jet energy profiles differentiate chiral- and vector-like DM fermions
- Chirality of DM fermion reflects mass generation mechanism
- Deepen our understanding of dark sector


Back-up slides

Dark photons

- Possible to have hidden $U(1)'$ gauge group---dark photons
- MeV-scale vector mediators charged under $U(1)'$ enhance DM annihilation rate to get sizable excess in positron flux, Pospelov, Ritz 2009
- Resolve discrepancy between measured and calculated muon anomalous magnetic moment, though other models can too
Pospelov 2009; Endo et al. 2012

Dark photon search experiments

D. Curtin, 1412.0018



For $m_{A'} \sim 1$ GeV, kinetic mixing $\epsilon \gtrsim 10^{-5}$, so that A' decays within a length of $\sim \mathcal{O}(1)$ mm.