

Light WIMP with Scalar Mediator Implications from Higgs Precision

Po-Yan Tseng (Kavli IPMU)

Collaborators:

Shigeki Matsumoto (Kavli IPMU)


Yue-Lin Sming Tsai (Academia Sinica, Taiwan)

[ArXiv: 1811.03292](https://arxiv.org/abs/1811.03292)

5th, Dark Matter, Dark Energy, and Matter-antimatter
Asymmetry, 29th Dec. 2018

Introduction



- Dark matter **relic abundance** is about **25%** of our Universe.



Planck Collaboration

Introduction


- Thermally produced DM: **Freeze-out** mechanism.
- Weakly interacting DM(**WIMP**), gives the correct DM relic abundance.



Dan Hooper: 0901.4090

Introduction

- Thermally produced DM: Freeze-out mechanism.
- Weakly interacting DM(**WIMP**), gives the correct DM relic abundance.



Dan Hooper: 0901.4090

Introduction

- ♦ Thermally produced DM: Freeze-out mechanism.
- ♦ Weakly interacting massive particle(**WIMP**), gives the correct DM **relic abundance** $\Omega_X = 25\%$.

$$\Omega_X \propto \frac{1}{\langle \sigma v \rangle} \sim \frac{m_X^2}{g_X^4}$$

J.L. Feng, J. Kumar: 0803.4196

- ♦ $g_{\text{weak}} \simeq 0.65$ and $m_{\text{weak}} \simeq \mathcal{O}(100) \text{ GeV} - 1 \text{ TeV}$, weak interaction. We called **WIMP DM**.

$$\langle \sigma v \rangle \simeq 3 \times 10^{-26} \text{ cm}^3 / \text{sec}$$

Introduction

- Thermally produced DM: Freeze-out mechanism.
- Weakly interacting massive particle(**WIMP**), gives the correct DM **relic abundance** $\Omega_X = 25\%$.

$$\Omega_X \propto \frac{1}{\langle \sigma v \rangle} \sim \frac{m_X^2}{g_X^4}$$



J.L. Feng, J. Kumar: 0803.4196

- $g_{\text{weak}} \simeq 0.65$ and $m_{\text{weak}} \simeq \mathcal{O}(100) \text{ GeV} - 1 \text{ TeV}$, weak interaction. We called **WIMP DM**.

$$\begin{aligned} 10^{-3} &\lesssim g_X \lesssim 3 \\ 10 \text{ MeV} &\lesssim m_X \lesssim 10 \text{ TeV} \end{aligned}$$

Introduction

- Supersymmetry theory predict the mass of WIMP around $O(100)$ GeV to 1 TeV.
- It is constrained from direct detection searches.




Towards Dark Matter Discovery 2018

P. Cushman et. al.:1310.8327v2

Introduction

- ♦ Sub-GeV thermal produced WIMP. **Light WIMP**.
- ♦ Lee-Weinberg limit. Require thermal DM mass larger than GeV.

B. W. Lee, S. Weinberg: PRL. 39(1977), 165.




- ♦ Constraint from Cosmic Microwave Background (**CMB**):
$$\langle \sigma v \rangle_{\text{CMB}} / m_\chi \lesssim 3 \times 10^{-28} \text{ cm}^3 \text{ s}^{-1} \text{ GeV}^{-1}$$
- ♦ Cannot give correct DM relic density, if DM lighter than **10 GeV**.

Introduction

- Sub-GeV thermal produced WIMP. **Light WIMP**.
- Lee-Weinberg limit. Require thermal DM mass larger than GeV.

B. W. Lee, S. Weinberg: PRL. 39(1977), 165.



- Constraint from Cosmic Microwave Background (**CMB**):


$$\langle \sigma v \rangle_{\text{CMB}} / m_\chi \lesssim 3 \times 10^{-28} \text{cm}^3 \text{s}^{-1} \text{GeV}^{-1}$$

- Cannot give correct DM relic density, if DM lighter than **10 GeV**.

$$\langle \sigma v \rangle \simeq 3 \times 10^{-26} \text{cm}^3 / \text{sec}$$


Introduction

- ◆ Relic abundance and DM annihilation.
- ◆ $DM+DM \rightarrow SM+SM$:



Introduction


- ◆ Relic abundance and DM annihilation.
- ◆ $DM+DM \rightarrow \text{mediator}+\text{mediator} \ \& \ \text{mediator} \rightarrow SM+SM$



- ◆ The mass of mediator is also Sub-GeV.

Introduction

- ◆ Relic abundance and DM annihilation.
- ◆ $DM+DM \rightarrow \text{mediator}+\text{mediator} \ \& \ \text{mediator} \rightarrow SM+SM$




DM annihilation and relic abundance 25%

- ◆ The mass of mediator is also Sub-GeV.

Introduction


- ◆ Relic abundance and DM annihilation.
- ◆ $DM+DM \rightarrow \text{mediator}+\text{mediator} \ \& \ \text{mediator} \rightarrow SM+SM$



- ◆ The mass of mediator is also Sub-GeV.

Introduction

- ◆ Relic abundance and DM annihilation.
- ◆ $\text{DM} + \text{DM} \rightarrow \text{mediator} + \text{mediator} \ \& \ \text{mediator} \rightarrow \text{SM} + \text{SM}$




p - wave : $\langle \sigma v \rangle \propto v^2$, avoid CMB constraint

- ◆ The mass of mediator is also Sub-GeV.

Introduction


- Thermal equilibrium.
- $WIMP \leftrightarrow \text{mediator} \leftrightarrow SM$.



S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Introduction





- Thermal equilibrium.
- $\text{WIMP} \leftrightarrow \text{mediator} \leftrightarrow \text{SM}$.



S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Introduction

- Thermal equilibrium.
- $WIMP \leftrightarrow \text{mediator} \leftrightarrow SM$.



S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Light WIMP with scalar mediator

- Minimal Model. Gauge invariant and renormalizability.
- Majorana DM and a scalar mediator:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2}\bar{\chi}(i\not{\partial} - m_{\chi})\chi + \frac{1}{2}(\partial\Phi)^2 - \frac{c_s}{2}\Phi\bar{\chi}\chi - \frac{c_p}{2}\Phi\bar{\chi}i\gamma_5\chi - V(\Phi, H),$$

$$V_H(H) = \mu_H^2 H^\dagger H + \frac{\lambda_H}{2}(H^\dagger H)^2,$$

$$V_\Phi(\Phi) = \mu_1^3 \Phi + \frac{\mu_\Phi^2}{2}\Phi^2 + \frac{\mu_3}{3!}\Phi^3 + \frac{\lambda_\Phi}{4!}\Phi^4,$$

$$V_{\Phi H}(\Phi, H) = A_{\Phi H}\Phi H^\dagger H + \frac{\lambda_{\Phi H}}{2}\Phi^2 H^\dagger H.$$

Light WIMP with scalar mediator

- Minimal Model. Gauge invariant and renormalizability.
- Majorana DM and a scalar mediator:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2}\bar{\chi}(i\not{\partial} - m_{\chi})\chi + \frac{1}{2}(\partial\Phi)^2 - \frac{c_s}{2}\Phi\bar{\chi}\chi - \frac{c_p}{2}\Phi\bar{\chi}i\gamma_5\chi - V(\Phi, H),$$

$$V_H(H) = \mu_H^2 H^\dagger H + \frac{\lambda_H}{2} (H^\dagger H)^2,$$

$$V_\Phi(\Phi) = \mu_1^3 \Phi + \frac{\mu_\Phi^2}{2} \Phi^2 + \frac{\mu_3}{3!} \Phi^3 + \frac{\lambda_\Phi}{4!} \Phi^4,$$

$$V_{\Phi H}(\Phi, H) = A_{\Phi H} \Phi H^\dagger H + \frac{\lambda_{\Phi H}}{2} \Phi^2 H^\dagger H.$$

Light WIMP with scalar mediator

- Minimal Model. Gauge invariant and renormalizability.
- Majorana DM and a scalar mediator:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2}\bar{\chi}(i\not{\partial} - m_{\chi})\chi + \frac{1}{2}(\partial\Phi)^2 - \frac{c_s}{2}\Phi\bar{\chi}\chi - \frac{c_p}{2}\Phi\bar{\chi}i\gamma_5\chi - V(\Phi, H),$$

$$V_H(H) = \mu_H^2 H^\dagger H + \frac{\lambda_H}{2}(H^\dagger H)^2,$$


$$V_\Phi(\Phi) = \mu_1^3 \Phi + \frac{\mu_\Phi^2}{2}\Phi^2 + \frac{\mu_3}{3!}\Phi^3 + \frac{\lambda_\Phi}{4!}\Phi^4,$$

$$V_{\Phi H}(\Phi, H) = \underline{A_{\Phi H}\Phi H^\dagger H} + \frac{\lambda_{\Phi H}}{2}\Phi^2 H^\dagger H.$$

Light WIMP with scalar mediator

- The $A_{\Phi H} \Phi H^\dagger H$ allowed the mixing between Higgs doublet and scalar singlet

$$\begin{pmatrix} h \\ \phi \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} h' \\ \phi' \end{pmatrix}$$



Light WIMP with scalar mediator

- Minimal Model. Gauge invariant and renormalizability.
- Majorana DM and a scalar mediator:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{2}\bar{\chi}(i\not{\partial} - m_{\chi})\chi + \frac{1}{2}(\partial\Phi)^2 - \frac{c_s}{2}\Phi\bar{\chi}\chi - \frac{c_p}{2}\Phi\bar{\chi}i\gamma_5\chi - V(\Phi, H),$$


$$V_H(H) = \mu_H^2 H^\dagger H + \frac{\lambda_H}{2}(H^\dagger H)^2,$$

$$V_\Phi(\Phi) = \mu_1^3 \Phi + \frac{\mu_\Phi^2}{2}\Phi^2 + \frac{\mu_3}{3!}\Phi^3 + \frac{\lambda_\Phi}{4!}\Phi^4,$$

$$V_{\Phi H}(\Phi, H) = A_{\Phi H}\Phi H^\dagger H + \frac{\lambda_{\Phi H}}{2}\Phi^2 H^\dagger H.$$

Light WIMP with scalar mediator

- The $\frac{\lambda_{\Phi H}}{2} \Phi^2 H^\dagger H$ give mediator-mediator-Higgs trilinear coupling.




Light WIMP with scalar mediator

- ♦ 7 parameters: $m_\chi, m_\phi, c_s, \sin \theta, \mu_\phi^2, \mu_3, \lambda_\Phi$

Light WIMP with scalar mediator

- 7 parameters: $m_\chi, m_\phi, c_s, \sin \theta, \mu_\phi^2, \mu_3, \lambda_\Phi$




Light WIMP with scalar mediator

- 7 parameters: $m_\chi, m_\phi, c_s, \sin \theta, \mu_\phi^2, \mu_3, \lambda_\Phi$



Light WIMP with scalar mediator

- 7 parameters: $m_\chi, m_\phi, c_s, \sin \theta, \mu_\phi^2, \mu_3, \lambda_\Phi$





Constraints

- ♦ Apriori constraints: **vacuum stability**.
- ♦ DM **relic density** & **Kinematic equilibrium** condition.
- ♦ Cosmology constraints: **BBN, Neff, CMB**.
- ♦ **Direct dark matter detection**.
- ♦ Collider constraints: **Kaon, B-meson, Higgs decay**.

Constraints

- ♦ Kinematic equilibrium condition+Kaon decay:



$$m_\chi \geq 10 \text{ MeV}$$



Constraints

- ◆ Kinematic equilibrium condition+Kaon decay:


$$m_\chi \geq 10 \text{ MeV}$$



Constraints

- Cosmology constraints: **BBN, Neff, CMB**. Assume mediator is thermal equilibrium with SM.

$$m_\phi \geq 6 \text{ MeV}$$




m_ϕ

BBN : hadronic decay $\tau_\phi < 1 \text{ second}$

Constraints

- Cosmology constraints: **BBN, Neff, CMB**. Assume mediator is thermal equilibrium with SM.


$$m_\phi \geq 6 \text{ MeV}$$



Constraints

- ♦ Cosmology constraints: **BBN, Neff, CMB**. Assume mediator is thermal equilibrium with SM.


$$m_\phi \geq 6 \text{ MeV}$$



Constraints

- Cosmology constraints: **BBN, Neff, CMB**. Assume mediator is thermal equilibrium with SM.

$$m_\phi \geq 6 \text{ MeV}$$




ϕ from relativistic to non-relativistic, inject entropy to $\gamma \Rightarrow$ change the $(T_D^{(\nu)}/T_D^{(\gamma)})$ and $(T_{\text{CMB}}^{(\nu)}/T_{\text{CMB}}^{(\gamma)})$

m_ϕ

$$\Delta N_{\text{eff}} : m_\phi \geq 6 \text{ MeV}$$


Results

- Under **present** constraints:




Results

- Under **future** constraints:




Light WIMP with scalar mediator

- 7 parameters: $m_\chi, m_\phi, c_s, \sin \theta, \mu_\phi^2, \mu_3, \lambda_\Phi$




Results

- From the allowed parameter space:



Results

- From the allowed parameter space:



Results

- ♦ Interesting signal from 125 GeV Higgs invisible decay at LHC.
- ♦ Because of the coupling $C_{\phi\phi h}$ the branching ratio of $h \rightarrow \phi\phi$ can be large.
- ♦ Current LHC limit is: $\text{Br}(h \rightarrow \text{invisible}) \lesssim 20\%$
- ♦ High luminosity LHC limit will be: $\text{Br}(h \rightarrow \text{invisible}) \lesssim 5\%$

Summary


- ◆ We demonstrated the **light WIMP DM** can be **Sub-GeV** with the help of **light scalar mediator** to maintain **thermal equilibrium** and give correct **relic density**.
- ◆ We wrote down a minimal model, which is gauge invariant and renormalizable.
- ◆ Many constraints are included.
- ◆ The 125 GeV Higgs decays into pair of long-live mediators as **invisible decay**. Can be searched at **LHC**.

Thank You !


Back Up

Introduction

◆ Dark matter



wikipedia.org Extended
rotation curve of M33



NASA




astro-ph/0504097




Hubble Space Telescope

Introduction

- ♦ Dark matter relic abundance is about 25% of our Universe.




Planck Collaboration



Wayne Hu: Department of Astronomy and Astrophysics U. of Chicago

Introduction

- Dark matter **relic abundance** is about **25%** of our Universe.




Introduction

- ◆ Properties of dark matter:
- ◆ I). Charge neutral.
- ◆ II). Stable or long-live.
- ◆ III). Gravity force.
- ◆ IV). Non-relativistic.
- ◆ V). 25% of our universe.

Introduction

- ◆ Properties of dark matter:
- ◆ I). Charge neutral.
- ◆ II). Stable or long-live.
- ◆ III). Gravity force.
- ◆ IV). Non-relativistic.
- ◆ V). 25% of our universe.



Wikipedia

Introduction

- ◆ Properties of dark matter:


- ◆ I). Charge neutral.

- ◆ II). Stable or long-live.

- ◆ III). Gravity force.

- ◆ IV). Non-relativistic.

- ◆ V). 25% of our universe.




2012
at
LHC

1897 by JJ
Thomson

Wikipedia

Introduction


- ◆ Properties of dark matter: **Beyond SM** particle.
- ◆ I). Charge neutral.
- ◆ II). Stable or long-live.
- ◆ III). Gravity force.
- ◆ IV). Non-relativistic.
- ◆ V). 25% of our universe.




Wikipedia

Introduction

- ◆ How to produced DM?
- ◆ Thermally produced: **Freeze-out** mechanism.




Wikipedia




$\langle \sigma v \rangle > \text{Hubble Expansion Rate}$
Thermal equilibrium

Introduction

- ◆ How to produced DM?
- ◆ Thermally produced: **Freeze-out** mechanism.



Wikipedia



$\langle \sigma v \rangle < \text{Hubble Expansion Rate}$
DM decouple

Introduction

- ♦ The mass of WIMP from O(1) MeV to O(100) TeV.
- ♦ O(100) TeV upper limit from the perturbation
- ♦ O(1) MeV lower limit from non-relativistic DM. Lighter DM freeze-out in relativistic.

J.L. Feng, J. Kumar: 0803.4196

$$\Omega_X \propto \frac{1}{\langle \sigma v \rangle} \sim \frac{m_X^2}{g_X^4}$$

$$\begin{aligned} 10^{-3} &\lesssim g_X \lesssim 3 \\ 10 \text{ MeV} &\lesssim m_X \lesssim 10 \text{ TeV} \end{aligned}$$

Introduction

- ♦ The mass of WIMP from O(1) MeV to O(100) TeV.
- ♦ O(100) TeV upper limit from the perturbative.
- ♦ O(1) MeV lower limit from non-relativistic DM. Lighter DM freeze-out in relativistic.


J.L. Feng, J. Kumar: 0803.4196

$$\Omega_X \propto \frac{1}{\langle \sigma v \rangle} \sim \frac{m_X^2}{g_X^4}$$

$$\begin{aligned} 10^{-3} &\lesssim g_X \lesssim 3 \\ 10 \text{ MeV} &\lesssim m_X \lesssim 10 \text{ TeV} \end{aligned}$$

Introduction



- ♦ Supersymmetry theory predict the mass of WIMP around $O(100)$ GeV to 1 TeV.
- ♦ It is constrained from direct detection searches.



KIPAC, Stanford U.

Constraints


- Relic abundance and thermal equilibrium.
- When $m_\chi \geq m_\phi$, t-channel WIMP annihilate into a pair of mediators.
- When $m_\chi \leq m_\phi$, s-channel WIMP annihilate into $m_\chi \sim m_\phi/2$. The resonance enhancement



Light WIMP with scalar mediator



- The $A_{\Phi H} \Phi H^\dagger H$ allowed the mixing between Higgs doublet and scalar singlet

$$\begin{pmatrix} h \\ \phi \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} h' \\ \phi' \end{pmatrix}$$




Results

- Under **present** constraints:




Results

Relic density




Results


$$\Delta N_{\text{eff}}$$



Results




Results




Kaon decay +
Kinematic equilibrium conduction

Results



Results





BBN (Big Band Nucleosynthesis)

$\tau_\phi < 1$ sec.



Results

- Under present constraints:






Results

- Under **future** constraints:




Introduction

- Thermal equilibrium.
- $WIMP \leftrightarrow \text{mediator} \leftrightarrow \text{SM}$.





S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Constraints



Constraints

WIMP \leftrightarrow mediator \leftrightarrow SM.




$$(m_\chi, c_s, m_\phi, \sin \theta, \mu_3) = (200\text{MeV}, 0.022, 100\text{MeV}, 10^{-6}, 10\text{MeV})$$

$$(200\text{MeV}, 0.1, 50\text{MeV}, 10^{-3}, 10\text{MeV})$$

Higgs Production at LHC

- ◆ The production mechanism: ggF, VBF, Vh, tth.



C.Grojean: 1708.00794


ProductionRate :

$$\text{ggF} = 80\%, \text{VBF} = 15\%, \text{Vh} = 4.5\%, \text{t}\bar{\text{t}}\text{h} = 0.5\%$$

K.Cheung, J.S.Lee, P.Y.Tseng: 1302.3794

Constraints

- When the mediator is lighter than 10 GeV. The LEP constraint is stronger than that from ILC, because of lower center mass energy. Y. Wang, J. List, M. Berggren: 1801.08164



ILC

- From the Higgs-mediator-mediator coupling, in small mixing angle limit, s.t. decay length is longer than $\sim 30\text{m}$. For example, $m_\phi = 20 \text{ GeV}, \sin \theta < 10^{-7}$

$$C_{\phi\phi h} \simeq \frac{2(m_\phi^2 - \mu_\Phi^2)}{v_H}$$
$$\Gamma(h \rightarrow \phi\phi) \simeq \frac{C_{\phi\phi h}^2}{32\pi m_h}$$

$$\Delta\text{BR}(h_{125} \rightarrow \text{invisible}) \lesssim 0.44\%$$

$$\Rightarrow C_{\phi\phi h} < 0.7 \text{ GeV}, \text{ or } |m_\phi^2 - \mu_\Phi^2| < 90 \text{ GeV}^2$$

Light WIMP with scalar mediator

- ♦ Higgs precision measurement at LHC:

$$\Delta\text{BR}(h_{125} \rightarrow ZZ) \lesssim 10\% \Rightarrow |\sin \theta| \lesssim 0.32$$



- ♦ At ILC (250GeV), improve the Higgs precision measurement:

$$\Delta\text{BR}(h_{125} \rightarrow ZZ) \lesssim 0.5\% \Rightarrow |\sin \theta| \lesssim 0.07$$

H. Baer et. al., ILC: 1306.6352

Light WIMP with scalar mediator

- Mediator width and branching ratio:




$$\sin \theta = 1, \quad \Gamma_{\phi} \propto \sin^2 \theta$$

S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Constraints

- Current experimental constraints for light WIMP. Lower mass limit for WIMP **9MeV**.




S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Constraints

- Current experimental constraints for light WIMP. Lower mass limit for WIMP **9MeV**.


thermal equilibrium



S.Matsumoto, Y.L Sming Tsai, P.Y. Tseng

Constraints


- Current experimental constraints for light mediator. Lower limit for mediator mass **1 MeV**.



S. Matsumoto, Y.L. Sming Tsai, P.Y. Tseng

Constraints

- Current experimental constraints for light mediator




The $\sin \theta$ too small,
but $C_{h\phi\phi}$ need to be large
to maintain thermal equilibrium.

S. Matsumoto, Y.L. Sming Tsai, P.Y. Tseng



Constraints

- Current experimental constraints for light mediator




Constraints

- ♦ Mediator produced at ILC



Constraints




- ◆ From the Higgs-mediator-mediator coupling

$$C_{\phi\phi h} \simeq \frac{2(m_\phi^2 - \mu_\Phi^2)}{v_H}$$

$$\Gamma(h \rightarrow \phi\phi) \simeq \frac{C_{\phi\phi h}^2}{32\pi m_h}$$

Constraints




- ◆ If the mixing angle with Higgs is very small, mediator becomes long-live particle.
- ◆ Invisible Higgs decay at ILC (250GeV):

$$\Delta\text{BR}(h_{125} \rightarrow \text{invisible}) \lesssim 0.44\%$$

H. Baer et. al., ILC: 1306.6352

Constraints

Invisible Higgs decay at ILC (250GeV):




S. Matsumoto, Y.L. Sming Tsai, P.Y. Tseng

Constraints

- Current experimental constraints for light mediator

$$C_{\phi\phi h} \simeq \frac{2(m_\phi^2 - \mu_\Phi^2)}{v_H}$$

$$\Gamma(h \rightarrow \phi\phi) \simeq \frac{C_{\phi\phi h}^2}{32\pi m_h}$$



$$\Delta\text{BR}(h_{125} \rightarrow \text{invisible}) \lesssim 0.44\%$$

H. Baer et. al., ILC:
1306.6352